

A MORPHOLOGICAL SURVEY ON THE REMARKABLE FLOWER OF *Dillenia indica* (DILLENiaceae)

Beatriz Dean Rizzo^{1,2}; Maria Juliana da Silva¹; Cláudio Gomes da Silva Jr Pedrosa³;
Geovana Gentilin Martins⁴; Michelle Rodrigues de Souza⁴; Luiz Antonio de Souza⁴;
Odair J. Garcia de Almeida¹ & J. Hugo Cota-Sánchez^{2*}

¹Universidade Estadual Paulista (UNESP), São Vicente, SP, Brasil

²University of Saskatchewan, Department of Biology, Saskatoon, SK, Canada

³Universidade Estadual de Maringá, PPG em Zootecnia, Maringá, PR, Brasil

⁴Universidade Estadual de Maringá, PPG em Biologia Comparada, Maringá, PR, Brasil

*Corresponding author: hugo.cota@usask.ca

Abstract

Within the flower, the stamens are among the structures that have undergone morphological and functional changes. The pantropical family Dilleniaceae includes ca. 400 species distributed in four subfamilies and 11 genera is among the flowering plants with flowers combining primitive and ancestral structures and poricidal anthers. The aim of this study was to illustrate the general structure of flowers and anthers in *Dillenia indica*. Flower parts are spirally organized including the apocarpous gynoecium composed of 16 to 22 carpels. The androecium has numerous free heterodynamous stamens, spirally organized in several concentric series that can be distinguished in peripheral and central series; the long anthers are poricidal. The ovary is epigynous and polycarpous with many ovules per carpel, axile placentation and anatropous ovules. The sepals persist during development and wrap an indehiscent fruit. All floral whorls are spirally arranged in a Magnoliid-fashion and showed some distinctive characters, such as perianth with persistent sepals that remain attached to the mature fruit; the androecium has at least two types of laminar poricidal stamens which shape is based on the locations (central versus peripheral), and the gynoecium is apocarpous. A mix of basal and derivative state of characters.

Keywords: Heteranthery, Micromorphology, Mosaic evolution, Poricidal anther

Resumo

Os estames estão entre as estruturas florais que sofreram alterações morfológicas e funcionais. Entre as plantas com flores com anteras poricidas, a família pantropical Dilleniaceae inclui cerca de 400 espécies distribuídas em quatro subfamílias e 11 gêneros. O objetivo desse estudo foi ilustrar a estrutura geral das flores, especialmente das anteras em *Dillenia indica*. As partes da flor são organizadas em espiral. O gineceu é apocárpico composto de 16 a 22 carpelos dispostos em espiral. O androceu possui numerosos estames livres, heterodínamos, organizados em espiral em várias séries concêntricas que podem ser distinguidas nas séries periférica e central; as anteras são poricidas. O ovário é epígino e policarpo com muitos óvulos por carpelo, placentação axial e óvulos anátropos. As sépalas persistem durante o desenvolvimento e envolvem o fruto indeiscente. Todos os verticilos florais são dispostos em espiral e apresentam algumas peculiaridades, como o perianto com sépalas persistentes que permanecem presas ao fruto maduro; o androceu tem pelo menos dois tipos de estames laminares poricidas, cuja forma é baseada na localização (central versus periférica), e o gineceu é apocárpico. Há, portanto, uma mistura de estados de caracteres basais e derivados.

Palavras-chave: Anteras poricidas, Evolução em mosaico, Heteranteria, Micromorfologia

Introduction

Approximately 135 million years ago, at the beginning of the Cretaceous, the first plants with specialized reproductive floral structures emerged (FRIIS et al., 2011). Ever since, numerous morphological traits have evolved in association with distinct forms of pollination and insects, often in a coevolutionary relationship favoring dispersal and reproductive success. Within the flower, the stamens are among the structures that have undergone morphological and functional changes. These reproductive organs typically consisting of anther, connective tissue and filament, carry pollen grains containing the male gametophyte. For pollen release to be effective, the anther walls are dehiscent and open so that pollen grains are released, attach to pollinator and are transferred onto the stigma of another flower. According to D'arcy (1996), the anther's functions include pollen production and long-distance dispersal. Stamen and anther shape and structure are

usually adapted to different floral pollination syndromes, thus diversifying the relationship between plants with their biotic pollinating agents.

Anthers can be diversified in several ways: shape, size, and type of dehiscence, which is the splitting of anther with subsequent release of pollen grains (ENDRESS, 1996). Most angiosperms have anthers with longitudinal or transverse slits (ENDRESS & STUMPF, 1991), but other irregular types of dehiscence include valvate and poricidal. Vogel (1978) described three phylogenetic types of flowers with distinct anther position, namely “*Magnolia* type”, “*Papaver* type”, and “*Solanum* type”, as a result of changes in flower shape and anthers over time. According to Bernhardt's (1996) the *Solanum* type is the most derived anther type and dehisces through apical pores, suggesting that poricidal anthers are one of the most advanced forms of anther dehiscence in angiosperms.

Among plant groups with poricidal anthers, the pantropical family Dilleniaceae includes ca. 400 species distributed in four subfamilies and 11 genera (STEVENS, 2019). Its members are characterized by several plesiomorphic characteristics including tall trees with solitary, multi-staminate flowers lacking nectaries, tricolporate pollen, apocarpous gynoecium, follicle or aggregate of follicles fruits (HORN, 2007). Within the family, the genus *Dillenia* L. includes 60 species bearing also basal morphological traits, such as, high number of spirally organized floral structures with apocarpous gynoecium, numerous carpels (16-22) and stamens (~650-850) and terminal idioblasts in leaves (character found in Magnoliaceae) (TUCKER, 1964). These characteristics led some authors to think that Dilleniaceae might be phylogenetically close to the ancestral angiosperm stock (STEBBINS, 1974). However, *D. indica* has flowers with poricidal anthers visited by bees which vibration facilitates the removal of pollen grains, i.e., buzz pollination (BUCHMANN, 1983). In addition, *D. indica* anthers have high amount of pollen and the stamens are arranged in characteristic central and peripheral position suggesting polymorphic heteranthery, a syndrome that involves more than one type of stamen with different function in the same flower (MÜLLER, 1883; VOGEL, 1978; VALLEJO-MARÍN et al., 2010). Most genera (64%) of the Dilleniaceae have anthers exclusively with longitudinal dehiscence, ca. 18% have anthers with longitudinal and poricidal anthers and ca. 18% of the genera have exclusively poricidal anthers (Table 1).

The flowers of *D. indica* are diurnal but remain open at night. The principal pollinators are bees of various sizes, which remove pollen from the anthers by buzz pollination (ENDRESS, 1997); no nocturnal floral visitors have been reported, but bats are the suspected pollinators in the original environmental species. The flowers have five

tepals and five sepals, but the stamens and carpels are numerous and spirally organized. The androecium is also organized in a spiral fashion. The gynoecium is apocarpous and forms a large multiple yellow fruit with acidic fragrance when ripe

The combination of both advanced and plesiomorphic features is an example of mosaic evolution in this family and may be result of a transformation of anther in relation to pollination selection. Few plant families display the existence of basal and advanced floral characters, which is relatively uncommon in angiosperms, lending this family a good study case for further investigation of floral characters. Within this context, in this study we aim to illustrate the general structure of flowers and anthers in *Dillenia indica*. The specific goals were: 1) to investigate the floral attributes and macro- and micromorphology of the anthers and pollen, and 2) to conduct a literature review to explore the phylogenetic occurrence of poricidal anthers within angiosperms to discuss *D. indica*'s floral characters in relation to the phylogenetic placement of the family within the IV APG scheme.

Material and methods

The study species – *D. indica* is a tree species naturally distributed from Madagascar and Sri Lanka to Australia and Fiji Islands, but introduced to Brazil as an ornamental tree (HOOGLAND, 1952).

Flowers were collected from four trees at the campus of the Maringá State University, Maringá, Paraná, Brazil, from February to March 2019. The identification of plant material was performed based on taxonomic keys and comparison with specimens cataloged in online databases such as World Checklist of Selected Plant Families (WCSP), The Plant List, Tropicos and the Missouri Botanical Garden. Permanent voucher specimens were deposited at the herbarium HUEM (Herbário da Universidade Estadual de Maringá). For morphological characterization *D. indica* flowers and reproductive structures were fixed in 50% EtOH, formaldehyde, PA 18: 1:1 acetic acid (FAA50) according to Johansen (1940) for at least 48 hours and then preserved in 50% EtOH for further analyses.

Measurement of floral characters – The number of tepals, sepals, carpels, stamen height, width and number were evaluated in ten fresh flowers collected from the four trees. For this, individual floral parts were dissected and arranged in progressive order for assessment. Measurements of height and width of the stamens and every floral structure

sampled were made using a digital caliper. The mean and standard deviation values were calculated with Microsoft Excel[®].

Macro- and micromorphology of anthers - The characterization of anthers in *D. indica* was performed using both stereoscope and scanning electron microscopy (SEM). Fresh samples were used for the former and for the latter the floral structures were fixed and preserved in 50% EtOH, then, dehydrated in a graded EtOH series: 70%, 80%, 90%, 100% and 100% for at least 1h on each step. An additional series of dehydration in EtOH/acetone was performed in diluted series as follows: 3:1 for 30 min, 1:1 for 30 min, 1:3 for 30 min, and 100% acetone for 30 min following Almeida et al. (2012). After dehydration the samples were critical point dried in liquid CO₂, sputter coated in gold and examined in SEM Phenom G2. The acquired images were edited using Photoshop CS3. For consistency, both peripheral and central anthers were characterized and described according to their morphological characteristics using at least three samples of each.

Literature review- In order to investigate the occurrence of poricidal anthers in angiosperms, an extensive literature review was performed based on journal articles and online manuscript databases, such as Google Scholar, University of Saskatchewan Library, Scopus, Web of Science, JSTOR, SciELO, Wiley Online Library, ScienceDirect, and Springer. The following key words as topics: poricidal anther, buzz pollination, angiosperms anther, anther, pore was invoked in this search. The anthers considered in this survey were all those that open through a pore, either single apical pore, two apical pores or subapical pores, as described by the authors. The outcome of this search was organized in a taxonomic table including class, family and genera (when available) of taxa reported with poricidal anthers (Table 4). A phylogenetic tree showing the orders with poricidal anthers angiosperms was generated using the online tool Interactive Tree of Life version 1.0 (iTOL[®]).

Results

Floral characters in Dillenia indica

The species is a tree up to 20 m (Fig. 1A) bearing solitary, terminal, and pendant (hanging as a bell) flowers subtended by a large peduncle developing in the apex of the branch (Fig. 1B). The blossoms are bisexual, actinomorphic and relatively large, measuring approximately 20cm, bowl-shaped and epigynous (Fig. 1C) with delicate

sweet-semi-acidic fragrance. Flower parts are spirally organized and bear a perianth with five flat, oval-oblong to curved petals white-cream and sepals spirally arranged (Figs. 1C, D; 2A, B). The perianth is composed of light green succulent oval sepals that range from 49.2mm to 57.9mm (54.5mm ave.) in length and 44.0mm to 52.6mm (48.0mm ave.) in width (Fig. 2A). The white caducous petals range from 65.0-91.0 mm (81.6mm ave.) in length and 38.0-53.0mm in width (47.4mm ave.) are also obovate to spatulate in shape (Fig. 2B) (Table 2). The androecium is also organized in spiral fashion and the stamens are arranged in a concentric peripheral and a central series (Figs. 2C-E). The stamens are distinct and the long anther forms a cylindrical-tubular structure (Figs. 2C, D). The gynoecium is apocarpous composed of 16 to 22 carpels (19 ave.) spirally arranged (Figs. 2F-I). The stigma lobes are white and smooth and vary from 16 to 22. The ovary is epigynous and polycarpous with many ovules per carpel, axile placentation and anatropous ovules. The sepals persist during development and wrap an indehiscent fruit (Figs. 2J-M).

Nocturnal observations were not performed, but during the day bees were observed, including *Scaptotrigona bipunctata* and *Apis mellifera scutellata*. Both bees collected only pollen grains during the day, but *S. bipunctata* was more active (Fig. 1C). Because the anthers are poricidal and hanging downwards, these bees have no apparent ability to promote buzz pollination or sonication. Thus, the bees use vibration and their jaws to damage the anthers and collect the pollen grains. According to Buchmann (1985) bees adapted to pollen collecting from poricidal anthers are in different families, e.g., Colletidae and Halictidae, among others.

Morphological characterization of androecium and stamens

The androecium has numerous yellow-creamy free stamens spirally organized in several concentric series that can be distinguished in peripheral and central series (Figs. 2C-E); hence, they are heterodynamous. That is, the stamens differ in shape and size and are inserted at different points at the base of the receptacle (Fig. 3A). The stamen insertion is basifixed (Figs. 3A-C). In both central and peripheral stamens, the filament texture has small wrinkles with square-rectangular cell shape on the epidermis (Figs. 3D, E). The central stamens are located between the stigmatic lobe spaces (Fig. 2C) surrounding the carpels and vary from 33 to 55 (ave. 47) in number, and measure on average 18.0 mm (16.5-23 mm) in length and 1.9 mm (1-2.45 mm) in width (Table 3). The number of the peripheral is larger than the central stamens and vary from 677 to 862 (ave. 762) and

measure on average 16.2 mm (ranging from 13.2-19.1 mm) in length and 1.0 mm (0.5-1.3 mm) in width.

Macro- and micromorphological characterization of anthers

The two types of anthers in *D. indica* flower are poricidal, with two thecas, and each theca has an apical pore, for a total of two apical pores/anther (Figs. 3A, B) forming a long cylindrical tube that involves nearly two thirds of the anther's length (Fig. 3A). Each of the anther's terminal pore has an operculum (Fig. 3C) that opens to release the pollen. The lateral region of the anther forms a continuous zig-zag pattern, mostly on the adaxial region (Fig. 3B). The anther epidermal surface has irregular-globose cells from the bottom to the top (Figs. 3D, F, G), which a rounded operculum opens on the adaxial surface facing the carpel (Fig. 3C). The filament cells in also irregular, with a rectangular shape (Figs. 3D, E). Both the central and peripheral anthers are typically loaded with small pollen grains of ca. 20-30µm in diameter (Figs. 4A-C). In central anthers, most pollen grains are spheric and tricolpate with pores and micropores of irregular shape on the surface (Figs. A, B). In peripheral anthers, the pollen is also spherical (Figs. 4C-F), along with some irregular pollen grains of oval (Fig. 4D), pentagonal (Fig. 4E), square (Fig. 4F) and other unusual irregular shapes was observed.

Phylogenetic occurrence of poricidal anthers in angiosperms

Poricidal anthers occur in 21 orders of flowering plants including monocots (Alismatales, Asparagales, Commelinales, Poales) and eudicotyledons (Apiales, Boraginales, Caryophyllales, Cucurbitales, Dilleniales, Ericales, Fabales, Gentianales, Lamiales, Malpighiales, Malvales, Myrtales, Oxalidales, Rosales, Santalales, Solanales and Zygophyllales) (Fig. 5).

According to APG's (2001) phylogenetic tree, poricidal anthers were found in four monocotyledonous orders, eight rosids, six asterids and three uncertain groups in the angiosperm phylogenetic tree, among them: Dilleniales, Santalales and Caryophyllales. In total, 50 families have at least one species with a poricidal anther, grouping 166 genera (Table 4). The order Ericales accounts 35% of the poricidal anther genera, followed by Malpighiales with 14%, accounting together almost 50% of poricidal anthers genera.

Discussion

The vegetative and morphological features of *D. indica* are quite characteristic and encompass an amalgamation of plesiomorphic and advanced characters. The large and scented hermaphrodite flower in particular has special attributes. All floral whorls are spirally arranged in a Magnoliid-fashion and showed some peculiarities, such as the perianth with persistent sepals that remain attached to the mature fruit; the androecium has at least two types of laminar poricidal stamens which shape is based on the locations (central versus peripheral), and the gynoecium is apocarpous. A mix of basal and derivative state of characters.

Stamen dimorphism and deviations from typical stamen morphology, including anther diversity, size, and shape, have intrigued scientists for many years. Darwin (1899) suspected that stamens related to heteranthery had different functions but was unable to understand their roles. Lloyd (2000) discussed the “feeding” anthers, which produces “feed pollen.” Usually, plants that have flowers with this kind of anther are nectarless and the main floral reward is pollen. The difference between central and peripheral anthers in *D. indica* offer the possibility of different pollen types, potentially a “feeding pollen,” that is pollen grains with the necessary proteins, fats and carbohydrates. Even with the lack in the studies regarding pollen viability, it is possible that in some peripheral anthers the pollen grains are “cheap” and not necessarily nutritious and/or not reproductively viable. Some pollen analyzes regarding Dilleniaceae, e.g., Furness (2007), Furness & Rudall (2004), reported that in *Tetracera* spp., pollen grains are inaperturate, just like some found in peripheral anthers of *D. indica*; however, it was not noticed any other morphological difference (size, shape or epidermal wall sculpturing) between triporate and inaperturate grains. Therefore, as Anderson & Symon (1989) discussed, it is probable that the peripheral anther’s pollen is a reward to pollinators, such as in some *Solanum* L. flowers. However, studies regarding the difference between introduced plants and native plants could be interesting to compare both peripheral anther’s pollens to confirm whether there is a difference between pollen in peripheral and central anther in the natural environmental. Also, a survey regarding both peripheral and central pollen’s viability would be interesting to better understand *D. indica*’s pollination system.

According to Vallejo-Marín et al. (2010), heteranthery is highly linked to poricidal anthers, whose mechanisms may be related to the reduction of pollen consumption by floral visitors, leaving more pollen for pollination. However, *A. mellifera* and *S. bipunctata* are not bees recognized for buzz pollination; nevertheless, small

disruptions in their surface were observed throughout the stamen, indicating a possible pollen “theft” for food functions. Endress (1997) observed many floral visitors in *D. indica*, *D. philippinensis* and *D. suffruticosa*, and *Xylocopa* sp. was considered an efficient pollinator. According to the author, these bees grab many poricidal anther stamens and vibrate their thorax, agitating the anthers and releasing the pollen grains. Then, the bee gathers another group of stamens and repeat the action so that it can transfer the pollen wedged in their body and to the stigmata lobe.

Although there are some floral visitors reports to *Dillenia* sp. flowers, there are almost no references regarding their pollinators. Corner (1988) mentions bees, small beetles and flies as likely pollinators. In general, its floral visitors are considered apiforms, among them *Xylocopa* sp., but often small bees, which are not probable pollinators (BURKILL, 1916; 1919) such as *Bombus morio* (AMARAL NETO, 2011) and *A. mellifera* (NOWICKE & MESELSON, 1984).

Because of the lightweight of pollen grains and the wide opening of the flower during anthesis, there is the possibility that anemophily might be the legitim pollen transport agent of specie. The evolution of anemophily is often explained regarding the environmental variation, caused by typical weather or the evolution of earth climate in the past million years (CULLEY et al., 2002). Although it is considered difficult for poricidal anthers to release pollen, *D. indica*'s pollen quantity is very large and can be released with a simple hand touch. In addition, according to Culley et al. (2002), *D. indica* has some traits typically associated with wind and biotic pollination, such as pollen size, number of pollen and absent nectary. It may indicate a different pollination syndrome depending on whether the specimens are in a native or introduced environment.

The first approach for the taxonomic revision of poricidal anthers was made by Harris (1905), whose apical dehiscence stamens were divided into 7 types. Later, Buchmann (1978) revived Harris's poricidal anthers classification, but with the addition of the buzz pollination, grouping a total of 72 families. However, the complete list of plants with poricidal terminal openings in Buchmann (1978) presents unconventional orders, such as Laurales, Canellales and Liliales, known for valvar or filamentous anthers. It happened because Buchmann considered poricidal anthers the species whose has short apical slits, short transverse slits, valves or legitim apical pores.

Acknowledgments

We thank R. Vera and D. Falconer for comments to improve the report. This survey was funded by grants from CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico – Brazil, PIBIC - No 53580/2019) and Emerging Leaders in the Americas Program (Canada Government). Also, we thank the personnel of the Universidade Estadual de Maringá for facilitating access to lab equipment, and the Plant Systematics Laboratory of the University of Saskatchewan and the Plant Morphology Lab of São Paulo State University (Unesp, Brazil) for all support and help to discuss and organize the results.

References

- AFFRE, L.; THOMPSON, J.D. Population genetic structure and levels of inbreeding depression in the Mediterranean island endemic *Cyclamen creticum* (Primulaceae). **Biological Journal of the Linnean Society**, v. 60, p.527-549, 1997.
- ALMEIDA, O.J.G.; PAOLI, A.A.S.; COTA-SÁNCHEZ, J.H. A macro- and micromorphological survey of floral and extrafloral nectaries in the epiphytic cactus *Rhipsalis teres* (Cactoideae: Rhipsalideae). **Flora**, v. 207, p. 119-125. 2012. <https://doi.org/10.1016/j.flora.2011.11.004>
- AMARAL, M.C.E.; BITTRICH, V. Ochnaceae. In: KUBITZKI, K. (ed.). **The Families and Genera of Vascular Plants - Flowering Plants. Eudicots**, 1st ed. Berlin, Springer-Verlag, p. 253-268, 2014.
- AMARAL NETO, L.P. **Forma e funcionamento das flores com quilha invertida em Faboidae e interações comportamentais com abelhas visitantes**. Dissertation, Universidade Federal do Paraná, Curitiba, Paraná, Brazil, 2011.
- ANDERSON, W.R. Notes on neotropical Malpighiaceae II. **Contributions from the University of Michigan Herbarium**, v. 16, p. 55-108, 1987.
- ANDERSON, G.J.; SYMON, D.E. Functional dioecy and andromonoecy in *Solanum*. **Evolution**, v. 43, p. 204-219, 1989.
- AYMARD, G.A.C. Dilleniaceae novae Neotropicae IX: *Neodillenia*, a new genus from the Amazon basin. **Harvard University Herbaria**, v. 1, p.121-131, 1997.
- BANO, R.; ABID, R.; QAISER, M. Anther types of the monocots within flora of Karachi. **Pakistan Journal of Botany**, v. 40, p. 1839-1849, 2008.

- BARCELONA, J.F.; CO, L.L.; BALETE, D.S.; BARTOLOME, N.A. *Rafflesia aurantia* (Rafflesiaceae): a new species from Northern Luzon, Philippines. **Gardens Bulletin Singapore**, v. 61, p. 17-27, 2009.
- BEARDSSELL, D.V.; O'BRIEN, S.P.; WILLIAMS, E.G.; KNOX, R.B.; CALDER, D.M. Reproductive biology of Australian Myrtaceae. **Australian Journal of Botany**, v. 5, p. 511-526, 1993.
- BERNHARDT, P. Anther adaptations in animal pollination. In: D'ARCY, W.G.; KEATING, R.C. (eds.). **The Anther: Form, Function and Phylogeny**, 1st ed. New York: Cambridge University Press, p. 192-220, 1996.
- BRUMMITT, R.K.; BANKS, H.; JOHNSON, M.A.T.; DOCHERTY, K.A.; JONES, K.; CHASE, M.W.; RUDALL, P.J. Taxonomy of Cyanastroideae (Tecophilaeaceae): a multidisciplinary approach. **Kew Bulletin**, v. 53, p. 769-803, 1988.
- BUCHMANN, S.L. **Vibratile ("Buzz") Pollination in Angiosperms with Poricidally Dehiscent Anthers**. PhD Dissertation, University of California, Davis, United States, 1978.
- BUCHMANN, S.L. Buzz pollination in angiosperms. In: JONES, C.E.; LITTLE, R.J. (eds.) **Handbook of Experimental Pollination Biology**, 1st ed. New York: Scientific and Academic Editions, p. 73-113, 1983.
- BUCHMANN, S.L. Bees use vibration to aid pollen collection from non-poricidal flowers. **Journal of the Kansas Entomological Society**, v. 58, p. 517-525, 1985.
- BURKILL, I.H. Notes on the pollination of flowers in India. - Note No. 8. Miscellanea. **Journal of the Royal Asiatic Society of Bengal**, v. 12, p. 239-265, 1916.
- BURKILL, I.H. Some notes on the pollination of flowers in the Botanic Gardens, Singapore, and in other parts of the Malay Peninsula. **Gardens' Bulletin Singapore**, v. 2, p.165-176, 1919.
- CAROLIN, R.C.; BITTRICH, V. Pittosporaceae. In: KUBITZKI K. (ed.). **The Families and Genera of Vascular Plants - Flowering Plants – Vol. XV - Eudicots**. Berlin: Springer, p. 539-547, 2018.
- CARTER, S. Revision of *Walleria* and *Cyanastrum* (Tecophilaeaceae). **Kew Bulletin**, v. 16, p. 185-195, 1962.
- CONRAN, J.G.; CAROLIN, R. Byblidaceae. In: KUBITZKI, K. (ed.). **The Families and Genera of Vascular Plants – Vol. VI Flowering Plants. Eudicots**, 1st ed. Berlin: Springer, p. 45-49, 2004.

- CORNER, E.J.H. **Wayside trees of Malaya**. Vol. 1. Kuala Lumpur: Malayan Nature Society, p. 296–304, 1988.
- CULLEY, T.M.; WELLER, S.G.; SAKAI, A.K. The evolution of wind pollination in angiosperms. **Trends in Ecology & Evolution**, v.17, p. 361-369, 2002.
- DARWIN, F. The botanical work of Darwin. **Annals of Botany**, v. 13, p. 9-19, 1899.
- DICKISON, W.C. A comparative morphological study in Dilleniaceae, VI. Stamens and young stem. **Journal of the Arnold Arboretum**, v.51, p. 403-422, 1970.
- DICKISON, W.C. Comparative morphological studies in Dilleniaceae, VII. Additional notes on *Acrotoma*. **Journal of the Arnold Arboretum**, v.52, p.319-333, 1971.
- DOWNING, T.L.; LADOGES, P.Y.; DURETTO, M.F. Trichome morphology provides phylogenetically informative characters for *Tremandra*, *Platytheca* and *Tetratheca* (former Tremandraceae). **Plant Systematics and Evolution**, v.271, p.199-221, 2008.
- ENDRESS, P.K. Diversity and evolutionary trends in angiosperm anthers. In: D'ARCY, W.G.; KEATING, R.C. (eds.). **The Anther: Form, Function and Phylogeny**, 1st ed. New York: Cambridge University Press, p. 92-110, 1996.
- ENDRESS, P.K. Relationships between floral organization, architecture, and pollination mode in *Dillenia* (Dilleniaceae). **Plant Systematics and Evolution**, v.206, p.99–118, 1997.
- ENDRESS, P.K.; STUMPF, S. The diversity of stamen structures in 'Lower' Rosidae (Rosales, Fabales, Proteales, Sapindales). **Botanical Journal of the Linnean Society**, v.107, p.217-293, 1991.
- FRENCH, J.C. Patterns of endothecial wall thickenings in Araceae: subfamilies Pothoideae and Monsteroideae. **American Journal of Botany**, v. 72, p. 472-486, 1985.
- FRIIS, E.M.; CRANE, P.R.; RAUNSGAARD, K. **Early flowers and angiosperm evolution**. Cambridge: Cambridge University Press, 2011.
- FURNESS, C.A. Why does some pollen lack apertures? A review of inaperturate pollen in eudicots. **Botanical Journal of the Linnean Society**, v.155, p. 29-48, 2007.
- FURNESS, C.A.; RUDALL, P.J. Pollen aperture evolution – a crucial factor for eudicot success? **Trends in Plant Science**, v.9, p.154-158, 2004.
- GRAYUM MH (1990) Evolution and phylogeny of the Araceae. **Annals of the Missouri Botanical Garden**, v.77, p.628-697.

- HARDER, L.D.; BARCLAY, M.R. The functional significance of poricidal anthers and buzz pollination: controlled pollen removal from *Dodecatheon*. **Functional Ecology**, v. 8, p. 509-517, 1994.
- HARRIS, J.A. The dehiscence of anther by apical pores. Missouri Botanical Garden **Annual Report**, v. 1905, p. 167-257, 1905.
- HARTER, B.; LEISTIKOW, C.; WILMS, W.; TRUYLIO, B.; ENGELS, W. Bees collecting pollen from flowers with poricidal anthers in a south Brazilian araucaria forest: a community study. **Journal of Apicultural Research**, v. 41, p. 9-16, 2015.
- HIERN, W.P. **A monograph of Ebenaceae**. Transactions of the Cambridge Philosophical Society, Cambridge, 1873.
- HOLSTEIN, N.; GOTTSCHLING, M. Flowers of *Halgania* (Ehretiaceae, Boraginales) are set up for being buzzed and the role of intertwining anther trichomes. **Flora**, v. 240, p.7-15, 2018.
- HOOGLAND, R.D. A revision of the genus *Dillenia*. **Blumea: Biodiversity, Evolution and Biogeography of Plants**, v.7, p. 1-145, 1952.
- HOOGLAND, R.D. The genus *Tetracera* (Dilleniaceae) in the Eastern Old World. **Herbarium Bogoriense**, v.2, p.185-224, 1953.
- HORN, J.W. Dilleniaceae. In: KUBITZKI, K. (ed.). **The Families and Genera of Vascular Plants: Volume IX Flowering Plants – Eudicots**, 1st ed. Berlin: Springer, p. 132-154, 2007.
- ISLAM, M.S.U.; ABID, R.; QAISER, M. Anther types of dicots within flora of Karachi, Pakistan. **Pakistan Journal of Botany**, v. 40, p. 33-41, 2008.
- JOHANSEN, D.A. **Plant microtechnique**. London: McGraw-Hill Book Company, 1940.
- JOHNSON-FULTON, S.B. WATSON, L.E. Phylogenetic systematics of Cochlospermaceae (Malvales) based on molecular and morphological evidence. **Systematic Botany**, v. 42, p. 2871-282, 2017.
- KEATING, R.C. The comparative morphology of the Cochlospermaceae. III. The flower and the pollen. **Annals of the Missouri Botanical Garden**, v. 59, p. 282-296, 1972.
- KELLER, B.T.; BREEDLOVE, D.E. Two new species of *Saurauia* (Actinidiaceae) from Mexico. **Systematic Botany**, v. 6, p. 65-73, 1981.
- KELLER, J.A.; HERENDEEN, P.S.; CRANE, P.R. Fossil flowers and fruits of the Actinidiaceae from the Campanian (Late Cretaceous) of Georgia. **American Journal of Botany**, v. 83, p.528-541, 1996.

- KNUDSEN, J.T.; OLESON, J.M. Buzz-pollination and patterns in sexual traits in North European Pyrolaceae. **American Journal of Botany**, v. 80, p. 900-913, 1993.
- KNUTH, P.; LOEW, E. **Handbuch der Blütenbiologie**. Leipzig: Engelmann, 1899.
- KUBITZKI, K. *Doliocarpus*, *Davilla* und verwandte Gattungen (Dilleniaceae). **Mitteilungen der Botanischen Staatssammlung München**, v. 9, p. 1-105, 1971.
- KUBITZKI, K. Sarraceniaceae. In: KUBITZKI, K. (ed.). **The Families and Genera of Vascular Plants – Volume VI Flowering Plants. Eudicots**, 1st ed. Berlin: Springer, p. 422-425, 2004.
- KUIJT J, HANSEN, B. Misodendraceae. In: KUBITZKI, K. (ed.). **The Families and Genera of Vascular Plants - Flowering Plants. Eudicots**, 1st ed. Berlin: Springer, p. 121-124, 2015a.
- KUIJT, J.; HANSEN, B. Santalaceae. In: KUBITZKI, K. (ed.). **The Families and Genera of Vascular Plants - Flowering Plants. Eudicots**, 1st ed. Berlin: Springer, p. 143-165, 2015b.
- LADD, P.G.; PARNELL, J.A.N.; THOMSON, G. Anther diversity and function in *Verticordia* DC (Myrtaceae). **Plant Systematics and Evolution**, v. 219, p. 79-97, 1999.
- LLOYD, D.G. The selection of social actions in families: III. Reproductively disabled individuals and organs. **Evolutionary Ecology Research**, v. 2, p. 29-40, 2000.
- LUNA, I.; OCHOTENA, H. Phylogenetic relationships of the genera of *Theaceae* based on morphology. **Cladistics**, v.20, p. 223-270, 2004.
- MALÉCOT, V.; NICKRENT, D.L.; BAAS, P.; Van Den OEVER, L.; LOBREAUCALLEN, D. A morphological cladistic analysis of Olacaceae. **Systematic Botany**, v. 29, p. 569-586, 2004.
- MESQUITA-NETO, J.N.; BLÜTHGEN, N.; SCHLINDWEIN, C. Flowers with poricidal anthers and their complex interaction networks—Disentangling legitimate pollinators and illegitimate visitors. **Functional Ecology**, v. 32, p. 2321-2332, 2018.
- MÜLLER, F. Two kinds of stamens with different functions in the same flower. **Nature**, v. 27, p. 364–365, 1883.
- NOWICKE, J.W.; MESELSON, M. Yellow rain - a palynological analysis. **Nature**, v.309, p. 205-206, 1984.

- POZNER, R.; COCUCCI, A. Floral structure, anther development, and pollen dispersal of *Halophytum ameghinoi* (Halophytaceae). **International Journal of Plant Sciences**, v. 167, p.1091-1098, 2006.
- SCHÖNENBERGER, J.; ENDRESS, P.K. Structure and development of the flowers in *Mendoncia*, *Pseudocalyx*, and *Thunbergia* (Acanthaceae) and their systematic implications. **International Journal of Plant Sciences**, v. 159, p.446-465, 1998.
- SCHÖNENBERGER, J.; Von BALTHAZAR, M.; TAKAHASHI, M.; XIAO, X.; CRANE, P.R.; HERENDEEN, P.S. *Glandulocalyx upatoiensis*, a fossil flower of Ericales (Actinidiaceae/Clethraceae) from the Late Cretaceous (Santonian) of Georgia, USA. **Annals of Botany**, v.109, p.921-936, 2012.
- SIMPSON, M.G. Phylogeny and classification of the Haemodoraceae. **Annals of the Missouri Botanical Garden**, v. 77, p.722-784, 1990.
- STÅHL, B.; ANDERBERG, A.A. Myrsinaceae. In: KUBITZKI, K. (ed.). **The Families and Genera of Vascular Plants: Volume VI Flowering Plants – Dicotyledons**. 1st ed. Berlin: Springer, p. 266-281, 2004.
- STEBBINS, G.L. **Flowering Plants – Evolution above the Species Level**. Cambridge: Harvard University Press, 1974.
- STEVENS, P.F. **Angiosperm Phylogeny Website**. Version 14. Access on 25th November 2019. <http://www.mobot.org/MOBOT/research/APweb>
- STEVENS, P.F.; LUTEYN, J.; OLIVER, E.G.H.; BELL, T.L.; BROWN, E.A.; CROWDEN, A.S.; GEORGE, G.J.; JORDAN, P.; LADD, K.; LEMSON, C.B.; MCLEAN, Y.; MENADUE, J.S.; PATE, H.M.; STACE, C.M.W. Ericaceae. In: KUBITZKI, K. (ed.). **The Families and Genera of Vascular Plants - Flowering Plants – Volume VI - Dicotyledons**. Berlin: Springer, p. 145-194, 2004.
- STEVENS, P.F.; WEITZMAN, L. Sladeniaceae. In: KUBITZKI, K. (ed.). **The Families and Genera of Vascular Plants - Flowering Plants – Volume VI - Dicotyledons**. Berlin: Springer, p. 431-433, 2004.
- STRUWE, L.; PRINGLE, J.S. Gentianaceae. In: KUBITZKI, K. (ed.). **The Families and Genera of Vascular Plants - Flowering Plants – Volume XV - Eudicots**. Berlin: Springer, p. 453-503, 2018.
- THOMAS, J.L. A monographic study of the Cyrillaceae. **Contributions from the Gray Herbarium of Harvard University**, v.186, p.1-114, 1960.
- TUCKER, S.C. The terminal idioblasts in magnoliaceous leaves. **American Journal of Botany**, v. 51, p. 1051-1062, 1964.

- TUCKER, S.C. Trends in evolution of floral ontogeny in *Cassia* sensu stricto, *Senna*, and *Chamaecrista* (Leguminosae: Caesalpinoideae: Cassieae: Cassiinae); a study in convergence. **American Journal of Botany**, v.83, p.687-711, 1996.
- VALLEJO-MARIN, M.; Da SILVA, E.M.; SARGENT, R.D.; BARRETT, S.C.H. Trait correlates and functional significance of heteranthery in flowering plants. **New Phytologist**, v.188, p. 418-425, 2010.
- VENKATESH, C.S. The special mode of dehiscence of anthers of *Polygala* and its significance in autogamy. **Bulletin of the Torrey Botanical Club**, v.83, p.19-26, 1956.
- VOGEL, S.T. Evolutionary Trends in Pollen Flowers. In: RICHARDS, A.J. (ed.) **The Pollination of Flowers by Insects**, 6th.ed. London: Academic Press, p. 89-96, 1978.
- VORONTSOVA, M.S.; HOFFMANN, P. A phylogenetic classification of tribe Poranthereae (Phyllanthaceae Euphorbiaceae sensu lato). **Kew Bulletin**, v. 63, p.41-59, 2008.
- WEBER, A. Gesneriaceae. In: KUBITZKI, K. (ed.) **The Families and Genera of Vascular Plants – Volume VI Flowering Plants. Eudicots**, 1st ed. Berlin: Springer-Verlag, p. 63-158, 2004.
- WILSON, P.G. Myrtaceae. In: KUBITZKI, K. (ed.). **The Families and Genera of Vascular Plants – Volume X Flowering Plants Eudicots** 1st ed. Berlin: Springer, p. 212-271, 2010.
- WILSON, P.G.; O'BRIEN, M.M.; GADEK, P.A.; QUINN, C.J. Myrtaceae revisited: a reassessment of infrafamilial groups. **American Journal of Botany**, v. 88, p. 2013-2025, 2001.
- XU, Z.; DENG, M. Cannabaceae. **Identification and Control of Common Weeds**, v. 2, p. 119-124, 2017.

Table 1. Type of anther dehiscence in Dilleniaceae. Taxonomic arrangement follows Horn (2007).

Subfamily	Genera and (#spp)	Anther dehiscence	Reference
Delimoioideae	<i>Tetracera</i> (50)	Longitudinal	Hoogland (1953)
Dillenioideae	<i>Acrotrema</i> (9)	Poricidal	Dickison (1971)
	<i>Dillenia</i> (65)	Poricidal and Longitudinal	Hoogland (1952)
	<i>Didesmandra</i> (1)	Longitudinal	Dickison (1970)
	<i>Schumacheria</i> (3)	Poricidal	Dickison (1970)
Doliocarpodeae	<i>Curatella</i> (1)	Longitudinal	Kubitzki (1971)
	<i>Davilla</i> (25)	Longitudinal	Kubitzki (1971)
	<i>Doliocarpus</i> (45)	Longitudinal	Kubitzki (1971)
	<i>Neodillenia</i> (2)	Longitudinal	Aymard (1997)
	<i>Pinzona</i> (1)	Longitudinal	Kubitzki (1971)
Hibbertioideae	<i>Hibbertia</i> (225)	Poricidal and Longitudinal	Dickison (1970)

Table 2. Floral morphological characteristics of *D. indica*. Perianth and gynoecium morphometrics data was made using the same flowers used in Table 3.

Whorl/Organ	Character	
Perianth		
Sepals	Shape and margin	Oval, entire
	Color	Light green
	Number	5
Petals	Shape and margin	Elongated, entire
	Color	White
	Number	5
Androecium		
Central stamen	Attachment	Basifixed
	Dehiscence	Poricidal
	Number	~47
	Height	~16mm
	Width	~1mm
Peripheral stamen	Type	Basifixed
	Dehiscence	Poricidal
	Number	~841
	Height	~19mm
	Width	~1.9mm
Gynoecium		
Carpel	Number	Multicarpelar (~19)
Locule	Number	Multilocular (~19)
Ovary	Position	Epigynous
	Placentation	Axilar
Ovule	Attachment	Anatropous
	Number	~50 per carpel (~950 in total)

Table 3. Androecial characters of *Dillenia indica*. Sample size (n = 10)

Flower Number	#Central stamens	Height central stamens (mm)	Width central stamens (mm)	#Peripheral stamens	Height peripheral stamens (mm)	Width peripheral stamens (mm)
1	33	23.0	1.0	710	15.0	0.5
2	33	23.0	1.0	706	15.0	0.5
3	55	17.5	1.9	862	13.1	1.2
4	54	16.5	2.1	790	15.9	0.8
5	49	16.7	2.4	850	15.8	0.7
6	50	18.9	2.1	650	17.9	1.0
7	48	16.7	2.1	760	16.7	1.0
8	53	18.9	2.2	834	16.0	1.2
9	44	17.6	2.2	677	19.0	1.2
10	47	18.0	1.7	781	15.5	1.3
\bar{x}	47(± 7.5)	18.7(± 2.3)	1.9(± 0.5)	762(± 70.4)	16.2(± 1.5)	1.0(± 0.3)

Table 4. Taxonomic occurrence of poricidal anthers in angiosperms

Order	Family	Genus	Reference
Alismatales	Araceae		Buchmann, 1983
		<i>Culcasia</i>	French, 1985
		<i>Lagenandra</i>	Grayum, 1990
		<i>Cryptocoryne</i>	Grayum, 1990
Asparagales	Tecophilaeaceae		Harris, 1905
		<i>Conanthera</i>	Harris, 1905
		<i>Cyanastrum</i>	Brummitt et al., 1998
		<i>Cyanella</i>	Harris, 1905
		<i>Tecophilea</i>	Harris, 1905
		<i>Walleria</i>	Carter, 1962
		<i>Zephyra</i>	Harris, 1905
Poales	Poaceae		Knuth & Loew 1899
		<i>Dichanthium</i>	Bano et al., 2008
	Rapateaceae		Harris, 1905
		<i>Cephalostemon</i>	Harris, 1905
		<i>Rapatea</i>	Harris, 1905
		<i>Saxofridericia</i>	Harris, 1905
		<i>Schoenocephalium</i>	Harris, 1905
		<i>Spathathus</i>	Harris, 1905
		<i>Stegolepis</i>	Harris, 1905
	Commelinales	Commelinaceae	
<i>Cartonema</i>			Harris, 1905
<i>Dichorisandra</i>			Harris, 1905
Haemodoraceae			Buchmann, 1983
		<i>Conanthera</i>	Simpson, 1990
		<i>Cyanella</i>	Simpson, 1990
		<i>Odontostomum</i>	Simpson, 1990
		<i>Tecophilaea</i>	Simpson, 1990
		<i>Zephyra</i>	Simpson, 1990
Dilleniales		Dilleniaceae	
	<i>Acrotrema</i>		Dickison, 1971
	<i>Dillenia</i>		Hoogland, 1952
	<i>Hibbertia</i>		Dickison, 1971
	<i>Schumacheria</i>		Dickison, 1971
Zygophyllales	Krameriaceae		Buchmann, 1983
		<i>Krameria</i>	Harris, 1905
Oxalidales	Elaeocarpaceae		Buchmann, 1983
		<i>Tetrateca</i>	Downing et al., 2008
		<i>Tremandra</i>	Downing et al., 2008
Malpighiales	Achariaceae		Harris, 1905 (Flacourtiaceae)
		<i>Kiggelaria</i>	Harris, 1905 (Flacourtiaceae)
	Euphorbiaceae		Buchmann, 1983
		<i>Euphorbia</i>	Islam et al., 2008
	Malpighiaceae		Vogel, 1978
		<i>Janusia</i>	Anderson, 1987
		<i>Tristellateia</i>	Anderson, 1987
	Ochnaceae		Harris, 1905;
		<i>Blastemanthus</i>	Buchmann, 1978
		<i>Brackenridgea</i>	Buchmann, 1978
		<i>Cespedesia</i>	Amaral & Bittrich, 2014
		<i>Elvasia</i>	Amaral & Bittrich, 2014
		<i>Euthemis</i>	Buchmann, 1978
<i>Fleurydora</i>		Amaral & Bittrich, 2014	
<i>Godoya</i>		Amaral & Bittrich, 2014	
<i>Krukoviella</i>		Amaral & Bittrich, 2014	
<i>Lavradia</i>		Buchmann, 1978	
<i>Lophira</i>		Amaral & Bittrich, 2014	

		<i>Luxemburgia</i>	Amaral & Bittrich, 2014
		<i>Ochna</i>	Buchmann, 1978
		<i>Perissocarpa</i>	Amaral & Bittrich, 2014
		<i>Poecilandra</i>	Buchmann, 1978
		<i>Rhytidanthera</i>	Amaral & Bittrich, 2014
		<i>Sauvagesia</i>	Buchmann, 1978
		<i>Ouratea</i>	Vogel, 1978
	Phyllanthaceae		Vorontsova & Hoffmann, 2008
		<i>Poranthera</i>	Vorontsova & Hoffmann, 2008
	Rafflesiaceae		Buchmann, 1983
		<i>Rafflesia</i>	Barcelona et al., 2009
Fabales	Leguminosae		Harris, 1905
		<i>Cassia</i>	Harris, 1905
		<i>Chamaecrista</i>	Tucker, 1996
		<i>Dicorynia</i>	Harris, 1905
		<i>Distemonanthus</i>	Harris, 1905
		<i>Koompassia</i>	Harris, 1905
		<i>Labichea</i>	Harris, 1905
		<i>Storckiella</i>	Harris, 1905
	Polygalaceae		Venkatesh, 1956;
		<i>Polygala</i>	Venkatesh, 1956;
Rosales	Cannabaceae		Knuth & Loew 1899
		<i>Humulus</i>	Xu & Deng, 2017
Cucurbitales	Begoniaceae		Harris, 1905
		<i>Begonia</i>	Harris, 1905
Myrtales	Melastomataceae		Harter et al., 2015;
		<i>Comolia</i>	Mesquita-Neto et al., 2018
		<i>Lavousiera</i>	Mesquita-Neto et al., 2018
		<i>Leandra</i>	Harter et al., 2015
		<i>Macairea</i>	Mesquita-Neto et al., 2018
		<i>Miconia</i>	Mesquita-Neto et al., 2018
		<i>Pterolepis</i>	Mesquita-Neto et al., 2018
		<i>Rhynchanthera</i>	Harter et al., 2015
		<i>Tibouchina</i>	Harter et al., 2015
	Myrtaceae		Buchmann, 1983
		<i>Babingtonia</i>	Wilson et al., 2001
		<i>Beaufortia</i>	Wilson, 2010
		<i>Darwinia</i>	Wilson et al., 2001
		<i>Eucalyptus</i>	Beardsell et al., 1993;
		<i>Regelia</i>	Wilson, 2010
		<i>Verticordia</i>	Ladd et al., 1999
Malvales	Cochlospermaceae		Buchmann, 1983
		<i>Amoreuxia</i>	Johnson-Fulton et al., 2017
		<i>Cochlosperinum</i>	Keating, 1972
	Dipterocarpaceae		Harris, 1905
		<i>Monoporandra</i>	Harris, 1905
		<i>Stemonoporus</i>	Harris, 1905
	Malvaceae		Harris, 1905
		<i>Guichenotia</i>	Harris, 1905
		<i>Lasiopetalum</i>	Harris, 1905
		<i>Thomasia</i>	Harris, 1905
Santalales	Misodendraceae		Buchmann, 1983
		<i>Misodendrum</i>	Kuijt & Hansen 2015a
	Olacaceae		Buchmann, 1983
		<i>Anacolosa</i>	Malécot et al., 2004
		<i>Aptandra</i>	Malécot et al., 2004
		<i>Cathedra</i>	Malécot et al., 2004
		<i>Chaunochiton</i>	Malécot et al., 2004
		<i>Ongokea</i>	Malécot et al., 2004
		<i>Phanerodiscus</i>	Malécot et al., 2004

	Santalaceae		Buchmann, 1983 (alterado)
		<i>Korthalsella</i>	Kuijt & Hansen (2015b)
Caryophyllales	Halophytaceae		Pozner & Cocucci, 2006
		<i>Halophytum</i>	Pozner & Cocucci, 2006
Ericales	<u>Actinidiaceae</u>		Buchmann, 1983
		<i>Actinida</i>	Keller et al., 1996
		<i>Clematoclethra</i>	Keller et al., 1996
		<i>Saurauia</i>	Keller & Breedlove, 1981
	Clethraceae		Buchmann, 1983
		<i>Clethra</i>	Schönenberger et al., 2012
	Cyrillaceae		Buchmann, 1983
		<i>Purdiaea</i>	Thomas, 1960
	Ebenaceae		Buchmann, 1983
		<i>Diospyros</i>	Hiern, 1873
	Ericaceae		Buchmann, 1978
		<i>Agapetes</i>	Buchmann, 1978
		<i>Agauria</i>	Buchmann, 1978
		<i>Andromeda</i>	Buchmann, 1978
		<i>Arbutus</i>	Buchmann, 1978
		<i>Arctostaphylos</i>	Buchmann, 1978
		<i>Bejaria</i>	Stevens et al., 2004
		<i>Ceratostema</i>	Stevens et al., 2004
		<i>Chamaedaphne</i>	Stevens et al., 2004
		<i>Costera</i>	Stevens et al., 2004
		<i>Demosthenesia</i>	Stevens et al., 2004
		<i>Dimorphanthera</i>	Stevens et al., 2004
		<i>Diogenesia</i>	Stevens et al., 2004
		<i>Disterigma</i>	Stevens et al., 2004
		<i>Gaylussacia</i>	Stevens et al., 2004
		<i>Kalmia</i>	Stevens et al., 2004
		<i>Monotropis</i>	Stevens et al., 2004
		<i>Oreanthes</i>	Stevens et al., 2004
		<i>Pellegrini</i>	Stevens et al., 2004
		<i>Rhododendron</i>	Buchmann, 1978
		<i>Rhodothamnus</i>	Buchmann, 1978
		<i>Satyria</i>	Stevens et al., 2004
		<i>Semiramisia</i>	Stevens et al., 2004
		<i>Symphysia</i>	Stevens et al., 2004
		<i>Vaccinium</i>	Buchmann, 1978
	Lecythidaceae		Buchmann, 1978
		<i>Allantoma</i>	Buchmann, 1978
		<i>Asteranthos</i>	Buchmann, 1978
		<i>Barringtonia</i>	Buchmann, 1978
		<i>Bertholletia</i>	Buchmann, 1978
		<i>Cariniana</i>	Buchmann, 1978
		<i>Corythophora</i>	Buchmann, 1978
		<i>Couratari</i>	Buchmann, 1978
		<i>Eschweilera</i>	Buchmann, 1978
		<i>Grias</i>	Buchmann, 1978
		<i>Gustavia</i>	Buchmann, 1978
		<i>Lecythis</i>	Buchmann, 1978
	Myrsinaceae		Harris, 1905
		<i>Ardisia</i>	Ståhl & Anderberg 2004
		<i>Cybianthus</i>	Ståhl & Anderberg 2004
		<i>Monoporus</i>	Ståhl & Anderberg 2004
		<i>Tapeinosperma</i>	Ståhl & Anderberg 2004
		<i>Yunckeria</i>	Ståhl & Anderberg 2004
	Pentaphylacaceae		Buchmann, 1983
		<i>Pentaphylax</i>	Van Steenis 1955
	Primulaceae		Buchmann, 1983

		<i>Cyclamen</i> <i>Dodecatheon</i>	Affre & Thompson 1997 Harder & Barclay, 1994
	Pyrolaceae	<i>Chimaphila</i> <i>Moneses</i> <i>Orthilia</i> <i>Pyrola</i>	Buchmann, 1978 Buchmann, 1978 Buchmann, 1978 Knudsen & Oleson, 1993 Buchmann, 1978
	Sarraceniaceae	<i>Heliamphora</i>	Keller et al., 1996 Kubitzki 2004
	Sladeniaceae	<i>Ficalhoa</i> <i>Sladenia</i>	Buchmann, 1983 Luna & Ochoterena, 2004 Stevens & Weitzman, 2004
Gentianales	Gentianaceae	<i>Cotylanthera</i> <i>Exacum</i> <i>Orphium</i>	Harris, 1905 Struwe & Pringle, 2018 Struwe & Pringle, 2018 Struwe & Pringle, 2018
	Loganiaceae	<i>Gardneria</i>	Harris, 1905 Harris, 1905
	Rubiaceae	<i>Argostemma</i> <i>Strumpfia</i>	Buchmann, 1983 Harris, 1905; Harris, 1905;
Solanales	Solanaceae	<i>Cyphomandra</i> <i>Solanum</i>	Harris, 1905 Harris, 1905 Harris, 1905
Boraginales	Boraginaceae	<i>Halgania</i>	Buchmann, 1983 Holstein & Gottschling, 2018
Lamiales	Acanthaceae	<i>Mendoncia</i> <i>Pseudocalyx</i>	Buchmann, 1983 Schönenberger & Endress, 1998 Schönenberger & Endress, 1998
	Byblidaceae	<i>Byblis</i>	Buchmann, 1983 Conran & Carolin (2004)
	Gesneriaceae	<i>Beccarinda</i> <i>Leptoboea</i>	Harder & Barclay, 1994 Weber 2004 Weber 2004
	Pedaliaceae	<i>Pedaliium</i>	Islam et al., 2008 Islam et al., 2008
Apiales	Pittosporaceae	<i>Billardiera</i> <i>Cheiranthra</i>	Buchmann, 1983 Harris, 1905 Carolin & Bittrich 2018




Figure 1. Representative features of *Dillenia indica*. A: Individual trees cultivated at the campus of the Universidade Estadual de Maringá, Brazil. B: Pendulous terminal flower attached to tree; C: Hermaphrodite flower in early anthesis with visiting *Scaptotrigona bipunctata*; D: Flower showing five petals in late anthesis. Scale bars A: 2m; B: 10cm; C: 3cm; D: 6cm.




Figure 2. Floral and fruit attributes in *Dillenia indica*. A: Oblong to ovate sepals; B: Obovate to spatulate petals; C: Longitudinal section of the flower showing parts of the perianth, androecium and gynoecium; D: Flower with perianth removed displaying androecium with peripheral (red outline) and central (blue outline) laminar stamens and gynoecium with multiple lobes in the stigma; E: Central (above) and peripheral (below) stamens; F: Close-up of stigma and lobes; G: Gynoecium showing ovary, style and stigma lobes; H: Cross section of multicarpelate ovary and receptacle; I: Longitudinal section of the ovary and ovules; J: Fruit in longitudinal section K: Fruit in longitudinal and cross section; L: Fruit with seeds; M: Developmental series of the indehiscent fruit. Ca: Carpel; CST: Central Stamen; Gy: Gynoecium; Lo: Lobe; Ov: Ovary; Ovu: Ovule; OW: Ovary Wall; Pe: Petal; PST: Peripheral stamen; R: Receptacle Sd: Seed; Se: Sepal; Stg: Stigma. Scale bars A: 5cm; B: 2cm; C: 1cm; D: 0.3cm; E: 3cm.




Figure 3. Macro- and micromorphology of stamens in *Dillenia indica*. A: Peripheral laminar stamen with poricidal anther and arrow pointing to the pore magnification; B: Lateral view of zig-zagged central stamen; C: Poricidal anther with operculum in central stamen; D: Pore opening of central anther; E-G: close-up of epidermal cells in anthers as indicated by circles in D. F: Anther cells; G: Cell texture detail. An: Anther; CT: Connective tissue; Fi: Filament; Po: Pore. Scale bars A: 740 μ m; B: 300 μ m; C: 120 μ m; D: 90 μ m; E: 55 μ m; F: 15 μ m; G: 5 μ m




Figure 4. Different pollen types in peripheral and central anthers of *Dillenia indica*. A-B. Tricolpate pollen from central anthers. C-F. Tricolpate pollen from peripheral anthers. Scale bars A: 5 μ m; B: 20 μ m; C: 15 μ m; D: 10 μ m; E: 15 μ m; F: 10 μ m




Figure 5. Phylogenetic distribution of poricidal anthers in angiosperms. Orders in bold characters and red lines indicated genera with poricidal anthers.